

Your Dosing Tracker

For IBRANCE® (palbociclib) + Aromatase Inhibitor*

My IBRANCE start date this cycle: _____

month | day | year

My IBRANCE dose: _____

Record your start date, and mark every day you take your therapy.

Here's your checklist:

- ☐ Record the day you start taking IBRANCE, and tell your healthcare team your start date.
.....
- ☐ Weeks 1-3, take IBRANCE + an aromatase inhibitor. Week 4, take an aromatase inhibitor only. Be sure to follow your dosing schedule exactly as your healthcare team prescribes.
.....
- ☐ Check off your daily doses.
.....
- ☐ Print out a new tracker every time you start a cycle and record any dosage changes.
.....
- ☐ Remember to refill your prescription on time and always follow your healthcare team's instructions.
.....

*Aromatase inhibitor is a class of therapy that includes letrozole, anastrozole, and exemestane.

DAY OF THE WEEK							
WEEK ONE IBRANCE + AROMATASE INHIBITOR							
WEEK TWO IBRANCE + AROMATASE INHIBITOR							
WEEK THREE IBRANCE + AROMATASE INHIBITOR							
WEEK FOUR AROMATASE INHIBITOR ONLY							

Please see Indication and Important Safety Information on page 2.
Click for the [Full Prescribing Information](#) and [Patient Information](#) or visit [IBRANCE.com](#).



Indication

IBRANCE® (palbociclib) 125 mg capsules and tablets are a prescription medicine used in adults to treat hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-) breast cancer that has spread to other parts of the body (metastatic) in combination with an aromatase inhibitor as the first hormonal based therapy. It is not known if IBRANCE is safe and effective in children.

Important Safety Information

IBRANCE may cause serious side effects, including:

Low white blood cell counts (neutropenia). Low white blood cell counts are very common when taking IBRANCE and may cause serious infections that can lead to death. Your healthcare provider should check your white blood cell counts before and during treatment.

If you develop low white blood cell counts during treatment with IBRANCE, your healthcare provider may stop your treatment, decrease your dose, or may tell you to wait to begin your treatment cycle. Tell your healthcare provider right away if you have signs and symptoms of low white blood cell counts or infections such as fever and chills.

Lung problems (pneumonitis). IBRANCE may cause severe inflammation of the lungs during treatment that can lead to death. Tell your healthcare provider right away if you have any new or worsening symptoms, including:

- chest pain
- cough with or without mucus
- trouble breathing or shortness of breath

Your healthcare provider may interrupt or stop treatment with IBRANCE completely if your symptoms are severe.

What should I tell my healthcare provider before taking IBRANCE?

Before taking IBRANCE, tell your healthcare provider about all of your medical conditions, including if you:

- have fever, chills, or any other signs or symptoms of infection.
- have liver or kidney problems.
- are pregnant or plan to become pregnant; IBRANCE can harm your unborn baby.
 - **Females** who are able to become pregnant should use effective birth control during treatment and for at least 3 weeks after the last dose of IBRANCE. Your healthcare provider may ask you to take a pregnancy test before you start treatment with IBRANCE.
 - **Males** with female partners who can become pregnant should use effective birth control during treatment with IBRANCE for at least 3 months after the last dose of IBRANCE.
- are breastfeeding or plan to breastfeed. It is not known if IBRANCE passes into your breast milk. Do not breastfeed during treatment with IBRANCE and for 3 weeks after the last dose.

The most common side effects of IBRANCE include:

- Low red blood cell counts and low platelet counts. Call your healthcare provider right away if you develop any of these symptoms during treatment:
 - dizziness
 - shortness of breath
 - weakness
 - bleeding or bruising more easily
 - nosebleeds
- infections
- tiredness
- nausea
- sore mouth
- abnormalities in liver blood tests
- diarrhea
- hair thinning or hair loss
- increased blood creatinine

IBRANCE may cause fertility problems in males. This may affect your ability to father a child. Talk to your healthcare provider about family planning options before starting IBRANCE if this is a concern for you.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of IBRANCE. Call your doctor for medical advice about side effects.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. IBRANCE and other medicines may affect each other, causing side effects.

Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

Do not drink grapefruit juice or eat grapefruit products while taking IBRANCE as they may increase the amount of IBRANCE in your blood.

Tell your healthcare provider if you start a new medicine. Take IBRANCE exactly as your healthcare provider tells you.

When IBRANCE is used in combination with an aromatase inhibitor or fulvestrant, also read the Patient Information for the prescribed products.

Please click for the IBRANCE [Full Prescribing Information](#) and [Patient Information](#) or visit [IBRANCE.com](https://www.pfizer.com/ibrance).

PP-IBR-USA-7142 © 2025 Pfizer Inc.
All rights reserved. August 2025

