

# Your Dosing Tracker

## For IBRANCE® (palbociclib) + Fulvestrant

My IBRANCE start date this cycle: \_\_\_\_\_  
 month | day | year

My IBRANCE dose: \_\_\_\_\_

Record your start date and mark every day you take your therapy.

### Here's your checklist:

- Record the day you start taking IBRANCE and tell your healthcare team your start date.  
 .....
- Weeks 1-3 (Days 1-21), take IBRANCE. Week 4 (Days 22-28), don't take IBRANCE. Be sure to follow your dosing schedule exactly as your healthcare team prescribes.  
 .....
- Check off your daily IBRANCE doses.  
 .....
- Record the days you receive your fulvestrant injections.  
 .....
- Print out a new tracker every time you start a cycle and record any dosage changes.  
 .....
- Remember to refill your IBRANCE prescription on time and always follow your healthcare team's instructions.

DAY OF THE WEEK							
<b>WEEK ONE</b> IBRANCE							
<b>WEEK TWO</b> IBRANCE							
<b>WEEK THREE</b> IBRANCE							
<b>WEEK FOUR</b>							
NO IBRANCE							

Please see Indication and Important Safety Information on page 2. Click for the full [Prescribing Information](#) and [Patient Information](#) or visit [IBRANCE.com](http://IBRANCE.com).



## Indication

IBRANCE® (palbociclib) is a prescription medicine used to treat hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-) breast cancer that has spread to other parts of the body (metastatic) in combination with fulvestrant in women with disease progression following hormonal therapy.

## Important Safety Information

**IBRANCE may cause serious side effects, including:**

**Low white blood cell counts (neutropenia).** Low white blood cell counts are very common when taking IBRANCE and may cause serious infections that can lead to death. Your doctor should check your white blood cell counts before and during treatment.

If you develop low white blood cell counts during treatment with IBRANCE, your doctor may stop your treatment, decrease your dose, or may tell you to wait to begin your treatment cycle. Tell your doctor right away if you have signs and symptoms of low white blood cell counts or infections such as fever and chills.

**Before you take IBRANCE, tell your doctor if you:**

- have fever, chills, or any other signs or symptoms of infection.
- have liver or kidney problems.
- have any other medical conditions.
- are pregnant or plan to become pregnant; IBRANCE can harm your unborn baby.
  - Females who are able to become pregnant and who take IBRANCE should use effective birth control during treatment and for at least 3 weeks after stopping IBRANCE.
  - Males who are taking IBRANCE and have female partners who can become pregnant should use effective birth control during treatment with IBRANCE and for 3 months after the final dose of IBRANCE.
- are breastfeeding or plan to breastfeed; it is not known if IBRANCE passes into your breast milk. You and your doctor should decide if you will take IBRANCE or breastfeed. You should not do both.

**Common side effects of IBRANCE include:**

- Low red blood cell counts and low platelet counts. Call your doctor right away if you develop any of these symptoms during treatment:
  - dizziness
  - shortness of breath
  - weakness
  - bleeding or bruising more easily
  - nosebleeds
- infections
- tiredness
- nausea
- sore mouth
- abnormalities in liver blood tests
- diarrhea
- hair thinning or hair loss
- vomiting
- rash
- loss of appetite

IBRANCE may cause fertility problems in males. This may affect your ability to father a child. Talk to your doctor if this is a concern for you.

These are not all of the possible side effects of IBRANCE. For more information, ask your doctor. Tell your doctor if you have any side effect that bothers you or does not go away.

Tell your doctor about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. IBRANCE and other medicines may affect each other, causing side effects.

Do not drink grapefruit juice or eat grapefruit products while taking IBRANCE as they may increase the amount of IBRANCE in your blood.

Tell your doctor if you start a new medicine. Take IBRANCE exactly as your doctor tells you.

If you take too much IBRANCE, call your doctor right away or go to the nearest hospital emergency room.

Please click for the full [Prescribing Information](#) and [Patient Information](#) or visit [IBRANCE.com](http://IBRANCE.com).

